

# Reducing inequalities in Life Expectancy in Haringey *Actions for the Haringey Strategic Partnership.*

November 2006

## Summary

Local Authorities and Primary Care Trusts have a responsibility for promoting the health and well being of their residents. Overall, people in Haringey are living longer healthier lives than they did 20 years ago. However, on average people in Haringey still die younger than in England as a whole, and there are substantial differences in health between neighbourhoods within the borough. For example, men born in one the most deprived wards can expect to die eight years before men born in one of the most affluent- a shocking statistic.

The purpose of the Haringey Life Expectancy Action Plan is to enable the Haringey Strategic Partnership to deliver priority actions to improve life expectancy and reduce health inequalities to meet the 2010 PSA health inequalities targets.

Improving health and reducing health inequalities is a key priority for Haringey. As a spearhead area Haringey is aiming to **reduce the gaps in life expectancy and infant mortality by at least 10% between Haringey and the population as a whole by 2010**. Partners are being monitored on delivery of the following targets, achievement of which will contribute significantly to reducing the gap;

- Reduce mortality rates from heart disease and stroke and related diseases by at least 40% in people under 75, with at least a 40% reduction in the inequalities gap between Haringey and the population as a whole.
- Reduce mortality rates from cancer by at least 20% in people under 75, with a reduction in the inequalities gap between Haringey and the population as a whole of at least 6%
- Reduce adult smoking rates to 21% or less with a reduction in prevalence among routine and manual groups to 26% or less
- Reduce mortality from suicide and undetermined injury by at least 20%
- Halting the year-on-year rise in obesity among children under 11 by 2010 in the context of a broader strategy to tackle obesity in the population as a whole.

The full programme of Public Service Agreement Floor Targets includes a number of other targets which impact on health inequalities, including improvements in employment rates, housing, community safety, and education.

Reducing the gap in life expectancy is the overarching target of the Well-Being Theme Board, supported by the Local Area Agreement target to reduce the gap in premature mortality rates between Haringey and England, and between deprived and more affluent parts of the borough. It is also reflected in the Haringey Sustainable Community Strategy.

The causes of inequalities in health are multiple and complex, genetic and biological differences accounting for a small proportion. The other influences on health are largely avoidable and are the result of differences in life circumstances, the choices we are able to make about how we live, and access to services.

This action plan is based on a detailed analysis of routine data on disease-specific mortality and socio economic data in Haringey. Key partners from the Haringey Well Being Theme Board planned and hosted an event for stakeholders in February 2006<sup>1</sup> to discuss potential priorities to address low life expectancy and health inequalities in the borough. These discussions were informed by detailed analysis of current evidence on local need and effectiveness of interventions provided by policy leads from across the partnership.

The consultation drew out a number of underlying themes that were considered in the development of this plan:

1. The advantages of improved integration between, and co-location of, health and social care and other services to disadvantaged communities. In particular, the need to join up outreach services better to meet the full range of needs that affect well-being.
2. That interventions should be targeted on the basis of need, addressing issues that are particular to specific black and minority ethnic communities, people with mental health problems or disabilities, and individuals that do not speak English or who are relatively new to Haringey. Services should work together to establish the best ways to target services to those in need, whether it be geographically by neighbourhood, or by care group, or through improved assessment processes.
3. The important role of voluntary and community organisations in reaching marginalised and socially excluded communities, and how this can be integrated more effectively into care pathways.
4. The importance of focusing on children and people in their middle years in reaching the life expectancy target. The HSP should aim to ensure that children have the best possible start in life to maximise their life chances, and improve access to health services for middle-aged individuals to ensure that effective interventions to prevent avoidable illness are utilised (eg secondary/tertiary prevention).

The action plan was drawn together based on the analysis leading up to and outcomes from the Healthier Haringey event, and a feasibility review and prioritisation of relevant actions. The full action plan is presented in Section 1, identifying risk factors for the main causes of premature death and inequalities in

health in Haringey that are amenable to change, and actions that should be taken forward by partners to address them. There are varying levels of evidence available to support the effectiveness of these interventions ranging from a sense of good practice, through national policy to strong evidence that the intervention would be effective in improving health and reducing health inequalities.

Section 2 provides a summary of the data and evidence on why reducing health inequalities in Haringey is a priority for all partners in the HSP. Additional information and copies of background papers from the Healthier Haringey event are available from [karen.dunn@haringey.nhs.uk](mailto:karen.dunn@haringey.nhs.uk) on request.

A number of actions emerge from this detailed plan because they are supported by strong evidence of effectiveness and local need, and are not currently being comprehensively addressed. These should be taken forward as a matter of priority by the HSP:

### **Smoking**

1. Offer stop-smoking advice as part of clinical assessment in surgical care pathways.
2. Prepare local businesses for implementation of smoke-free legislation.
3. Expand coverage of the Haringey smoke-free award amongst venues serving deprived communities in Haringey, and amongst partner-accredited schemes such as child minder certification.

### **Physical activity**

4. Train primary health workers to identify inactive adults opportunistically, and provide advice on physical activity.
5. Expand opportunities for people to be physically active through walking and cycling, and access to sport, leisure and open spaces.
6. Expand targeted approaches to promoting physical activity (eg exercise referral schemes or volunteer walks) based on the outcomes of local and other evaluation.

### **Diet and nutrition**

7. Ensure all school achieve healthy school status accreditation, and that the food they provide meets national nutritional standards for school food.
8. Review the Haringey Food and Nutrition strategy focusing on groups with high levels of need eg people living on low incomes, and those living with cardiovascular disease, diabetes and cancer.
9. Develop a strategy to prevent obesity amongst adults and children, including care pathways.

### **Access to health services**

10. Develop needs-based approaches to commission primary care services, building on an equity audit of resource allocation to GP practices.
11. Ensure that prescription of statins to individuals with cardiovascular disease, or who have a greater than 20% risk of developing it over the next 10 years, is equitable.
12. Increase the proportion of GP practices with PCT-validated registers of patients with Coronary Heart Disease.

13. Ensure equitable implementation of NICE guidelines on hypertension and management of heart failure.
14. Increase uptake rates for cervical and breast screening, including non English-speaking communities.

### **Accidents**

15. Develop safer routes to school, and traffic safety measures.
16. Ensure that housing interventions include accident prevention measures such as fire safety, and removing the causes of trips and falls.

### **Suicide**

17. Develop a suicide prevention strategy incorporating mental health promotion, risk reduction amongst key population groups, and reducing the availability of suicide methods.

### **Infant mortality**

18. Develop a strategy to reduce the number of women booking late in their pregnancy for ante-natal care.
19. Establish systems to monitor the smoking status of, and interventions received by, families with children.
20. Develop smoking cessation services as a core element of care pathways developed within children's centres.
21. Develop a breastfeeding maintenance monitoring system using the child health surveillance system (6-8 week check), and use this to target interventions for women/families less likely to maintain breastfeeding.

### **Homes**

22. Develop housing condition assessment criteria and referral pathways to housing/environmental health services for use by a range of service providers visiting vulnerable people in their own homes.
23. Develop strategies to reduce fuel poverty and improve thermal comfort, particularly for households vulnerable to poor health.
24. Improve housing conditions in the private rented sector through the private sector housing service.

### **Employment**

25. Develop employment opportunities for disadvantaged groups, including people with mental health problems, with physical or learning disabilities, lone parents, and refugees.
26. Ensure Haringey residents have equitable access to the employment opportunities offered by local developments (eg Tottenham Hale) and our location in the London-Stanstead-Cambridge-Peterborough corridor Growth Area.
27. Evaluate the effectiveness of providing employment and income advice in GP practices to support individuals on incapacity benefit who want to return to work.

## **Education**

28. Support schools in developing provision that raises the achievement of pupils from Black and Minority Ethnic communities that are currently not achieving as well as the general population.
29. Ensure that all schools attain accreditation as meeting the national Healthy Schools standards.

This document will be presented to the five theme boards of the HSP for discussion and to agree a commissioning and monitoring framework for implementation. This will be overseen by the Well-Being Theme Board, and championed by the Director of Public Health Dr Ann-Marie Connolly.

### **For further details contact:**

Vicky Hobart  
Haringey TPCT: Public Health  
Vicky.hobart@haringey.nhs.uk

Helena Pugh  
Haringey Council: Social Services  
Helena.pugh@haringey.gov.uk

# Contents

<b>Section 1:</b>	What are the actions that the Haringey Strategic Partnership should take to improve life expectancy and reduce inequalities?	6
1.	smoking	7
2.	physical activity	9
3.	food and nutrition	11
4.	cardiovascular disease	13
5.	cancer	15
6.	accidents	16
7.	suicide	17
8.	access to health services	18
9.	infant mortality	19
10.	housing	20
11.	employment	22
12.	education	24
<b>Section 2:</b>	The case for action by the Haringey Strategic Partnership	25

## Section 1: What are the actions that the Haringey Strategic Partnership should take to improve life expectancy and reduce inequalities?

### 1. SMOKING

#### Objective: (inc. PSA & local targets)

**DH PSA3 / DfES PSA3:** Reduce adult smoking rates to 21% or less with a reduction in prevalence among routine and manual groups to 26% or less

**LAA target:** tbc

#### Current situation

Recent surveys/modelling from the HDA suggest Haringey is likely to have a smoking prevalence of 27-32%<sup>2</sup>. There are no local data on trends in smoking prevalence. However, national data show a reduction in overall prevalence of smoking over the past 30 years, with little change in smoking rates among those living on low incomes and those who are least advantaged<sup>3</sup>.

#### Initiatives To Reduce The Prevalence Of Smoking

Action	Target group	Evidence of effectiveness	Estimated cost	Delivery lead
Expansion of coverage of Haringey Smoke Free Award with focus on: <ul style="list-style-type: none"> <li>• targeting venues in east of borough</li> <li>• partnership-organisation accredited schemes e.g. child minder certification</li> </ul>	Venues in the east of the borough & accredited scheme users	Strong (4% reduction in workforce quitting <sup>4</sup> )	£7K 2006/07	E&H SSS
Preparation of local businesses for implementation of smoke free elements of Health Improvement and Protection Bill.	Local businesses likely to have high smoking prevalence	Strong (4% reduction in workforce quitting)	N/A	Environmental Health (LBH) Public Health (TPCT)
Make no-smoking policies a requirement when local NHS organisations and Haringey Council are contracting/commissioning	Commissioned service users	Good practice	N/A	Service Commissioners
Ensure that all strategic partners (e.g. police force, fire brigade and voluntary sector organisations) have policies in place to promote smoke-free messages	Strategic partners	Strong (4% reduction in workforce quitting)	N/A	Haringey Strategic Partnership
Increased enforcement of regulations on tobacco smuggling	Targeting should be based on assessment	Limited evidence on effectiveness of local measures	N/A	Environmental Health (LBH)

## Stop Smoking Initiatives

Action	Target group	Evidence of effectiveness	Estimated cost	Delivery lead
Continue development of NHS smoking cessation services: <ul style="list-style-type: none"> <li>▪ Establish choose and book system through GP practices from 2006.</li> <li>▪ Move level 3 clinic from NMH to Tynemouth Road</li> <li>▪ Establish level 3 clinic in Wood Green Library</li> <li>▪ Expand services in deprived parts of the borough</li> </ul>	Smokers, particularly in deprived areas	Strong. Cost per QALY £135 - £6472 <sup>5</sup>	N/A	E&H SSS
Offer of stop smoking advice as part of clinical assessment in surgical care pathways	Smokers awaiting elective surgery (about 5,739/yr)	Strong Estimated 433-904 elective patients would give up smoking, with a reduction in post-op complications of 77-160 <sup>6</sup>	Estimated annual cost saving due to reduction in complications and bed days of about £850,000	HTPCT to address through surgical care pathways
Maintain level 2 quit Smoking Programme for Haringey Council Staff	LBH staff	Strong	N/A	E&HSSS



## 2. PHYSICAL ACTIVITY

### Objective: (inc. PSA & local targets)

DCMS PSA3 By 2008 increase the number who participate in active sports at least 12 times a year by 3% and increase the number who engage in at least 30 minutes of moderate intensity level sport at least 3 times a week by 3%. A year-on-year incremental increase by 1% per annum in physical activity levels of the whole population (Choosing Health delivery recommendation). Physical activity also contributes to the PSA targets on CHD, cancer and obesity (halting the year-on-year increase in obesity amongst children under 11 by 2010, in the context of a broader strategy to tackle obesity in the population as a whole)

**LAA target:** tbc

### Current situation

On the basis of national data, it is estimated that in Haringey approx 123,000 adults <sup>7</sup> and 6,000 boys and 8,000 girls aged 2-15 are insufficiently active <sup>8</sup>. It is further estimated that of approximately 252 CHD deaths per year in Haringey, approx 94 are attributable to physical inactivity <sup>9</sup>.

Action	Target group	Evidence of effectiveness	Estimated cost	Delivery lead
Primary care health workers to be trained in opportunistic identification of inactive adults (using validated tool e.g. a GPPAQ), and advice to aim for 30 minutes of moderate activity on 5 days of the week (or more)	Inactive adults	Strong for giving advice (£750 to £3150 per QALY)	N/A	HTPCT Public Health
School Sport Co-ordinators to ensure that 5-16 year olds in Haringey engage in a minimum of two hours of high quality PE and school sport every week and that as many children as possible benefit from high quality play opportunities.	School children	National policy	N/A	Healthy Schools Programme
Train frontline staff to provide advice on physical activity including, practice nurses, Haringey Council Leisure centre staff, dieticians, physiotherapists, health care assistants.	Service users	Good evidence of effectiveness of primary care practitioners providing physical activity advice.	Approx £2,000 per course (20-27 participants)	Spearhead PCT Obesity Training Fund

<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Estimated cost</b>	<b>Delivery lead</b>
Promote access to open spaces by addressing safety concerns (e.g. through the provision of wardens, parks officers, improved lighting, community facilities).	Adults and Children	Good practice	N/A	LBH Environmental Services
Develop opportunities to promote physically active modes of transport e.g. walking and cycling.	Adults and Children	Good practice	N/A	LBH Environmental services
Exercise referral scheme being developed and evaluated as part of a randomised controlled trial in 3 deprived neighbourhoods in Northumberland Park, Bruce Grove and Noel Park wards.	Inactive Adults in 3 deprived neighbourhoods	To be established as part of RCT as recommended by NICE	N/A	NRF funding
Evaluate Haringey Get Up and Walk programme providing training for volunteer walk leaders to lead walks in their local communities	Inactive Adults	Insufficient-should only be conducted as part of a research study <sup>10</sup>	N/A	HTPCT Public Health
Evaluate Fit for Life Programme: 8-10 week courses of physical activity and healthy lifestyle advice for people at risk of CHD.	People at risk of CHD	To be evaluated	Approx £11,000 total (8 courses per year)	HTPCT Public Health
Evaluate Health for Haringey, a 5-year programme providing exercise and social support opportunities to 3,000 people in deprived areas	Physically inactive individuals in deprived areas	To be evaluated	£1 million over 5 years	Health for Haringey Programme (Big Lottery Fund)
Evaluate HPCT and LBH Health at Work programmes: promoting physical activity for employees of the PCT and LBH	Employees of the HPCT and LBH	To be evaluated	Approx £800 to-date	HTPCT-Public Health
Expand joint work between HTPCT and LBH to increase opportunities for physical activity for older people e.g. chair-based exercise sessions at Leisure Centres.	Older people	Good practice	N/A	Age Concern

### 3.FOOD and NUTRITION

**Objective: (inc. PSA & local targets)**

Halt the year on year rise in obesity among children under 11 by 2010 in the context of a broader strategy to tackle obesity in the population as a whole.

Also contributes to CHD and cancer PSA targets

**LAA target:** tbc

**Current situation**

There are no local data on obesity and food consumption. Nationally 22% of men and 23% of women in England are now obese, and has been trebling since the 1980s, and 70% of men and 63% of women are either overweight or obese. The greatest problems are in the lowest socioeconomic groups and amongst children and young people. Around 16% of 2 to 15 year olds are now obese.

Action	Target group	Evidence of effectiveness	Delivery lead
Strengthen implementation of infant feeding guidelines, including promotion of breastfeeding.	Parents of babies	Strong <sup>11</sup>	Children's service
Healthy Schools Programme to ensure all schools meet national standards for school food.	School children	National policy	Healthy Schools Programme
Develop children's access to healthy food through the extended schools programme e.g. breakfast clubs, particularly in areas of high deprivation.	School children in deprived areas	Good practice	Children's service
Establish baseline data on the prevalence of childhood obesity amongst reception and year 6 children Haringey, and systems for monitoring and acting on future trends.	School age children	National Policy	Children's Service and HTPCT Public Health
Update the Haringey Food and Nutrition Strategy focusing on those most in need particularly people living on low incomes and the those living with CHD, strokes, diabetes and cancer	Low income & people with CHD, stroke, diabetes and cancer	Good practice	HTPCT Public Health
Develop an obesity strategy and care pathway	People at risk of / with obesity	National policy	HTPCT LBH

<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Delivery lead</b>
Set standards and use contracting to improve the nutritional quality of meals provided by catering contractors e.g. in residential settings, day centres, meals on wheels, staff canteens and vending machines	Residents of residential settings	Good practice	HTPCT and LBH commissioners
Work with local businesses/suppliers to promote access to affordable healthy food (e.g. through positive award schemes)	Local population	Good practice	LBH Environmental Health
Work with local residents to share good practice in local food schemes e.g. allotments, food co-ops, community cafes, window boxes,	Local community groups	Good practice	HAVCO/HARCEN
Limit the number and density of fast food outlets	Consumers of fast food	Good practice	Environmental services
Target vulnerable and disadvantaged communities through community initiatives such as community nutrition assistants, and distribution of healthy eating messages through libraries etc	Disadvantaged communities	Good practice	HTPCT teaching programme, HAVCO, & HARCEN
Education/training programmes for service providers including school nurses to provide support and advice to prevent obesity and promote healthier eating	Service providers	Good practice	HTPCT Public Health

## 4.CARDIOVASCULAR DISEASE

### Objective: (inc. PSA & local targets) DH PSA1

Reduce mortality rates from heart disease and stroke and related diseases by at least 40% in people under 75, with at least a 40% reduction in the inequalities gap between Haringey and the population as a whole.

### Current situation

Haringey's cardiovascular disease mortality rate has fallen significantly from 152.6 per 100,000 population under 75 (152.6/100,000) in 1996/98 to 128.6/100,000 in 2002/04. However, the gap between the Haringey and England average widened by 14.7/100,000 over the same period to reach 31.9/100,000 in 2002/04<sup>12</sup>. In addition there are significant inequalities across the borough with mortality rates from CHD in those under 75 in Bruce Grove in 2000-4 89% higher than the national average<sup>13</sup>. Based on current trends, the LHO predicts that CHD mortality will fall by about 48% (from the 1995-7 baseline until 2010) but the gap in CHD mortality rates between Haringey and England will continue to increase.<sup>14</sup>

### PRIMARY PREVENTION

See Sections on Smoking, Physical Activity, Food, Employment And Education

### SECONDARY PREVENTION

Action	Target group	Evidence of effectiveness	Delivery lead
Increase percentage of GP practices with the following PCT-validated CHD registers: <ul style="list-style-type: none"> <li>asymptomatic patients with CHD risk &gt;30% over 10 years (PSA01b target)</li> <li>patients with CHD</li> <li>patients on CHD registers whose last measured cholesterol (measured within last 15 months) is 5mmol/l or less (PSA01d)</li> </ul>	Patients with CHD or at high CHD risk	Strong <sup>15</sup>	General practice / HTPCT Primary Care Performance
Prescription of statins to adults with clinical evidence of CVD and adults without CVD who have a >20% risk of developing CVD within 10 years	Patients at high risk of CVD & patients with CVD	Strong <sup>16</sup>	General Practice and HTPCT Pharmacy lead
Improving equity of access to health services (see section on ACCESS TO HEALTH SERVICES)			

**TERTIARY PREVENTION** (Treatment & Rehabilitation)

<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Delivery lead</b>
Implementation of PCT hypertension guidelines (in line with NICE guidelines)	Patients with hypertension	Strong <sup>17</sup>	HTPCT Public Health
Improve management of heart failure in line with NICE guidelines	Patients with heart failure	Strong <sup>18</sup>	HTPCT Public Health
Phase IV Community-based Cardiac rehabilitation group exercise programme	Adults with established CHD	Strong <sup>19</sup>	Participant contributions & HTPCT Public Health
Improve % of patients with heart attack who receive thrombolysis within 60 minutes	Patients with heart attack	Strong <sup>20</sup>	Whittington and NMUH.

## 5.CANCER

### Objective: (inc. PSA & local targets)

**DH PSA1** Reduce mortality rates from cancer by at least 20% in people under 75, with a reduction in the inequalities gap between Haringey and the population as a whole of at least 6%

### Current situation

Haringey's cancer mortality rate has fallen from 133.6 per 100,000 population under 75 (133.6/100,000) in 1996/98 to 124.0/100,000 in 2002/04. However, the England average has fallen faster over the same period. Haringey's cancer mortality rate is now marginally 4% above the England average, and the gap between the two beginning to widen<sup>21</sup> Based on current trends, the LHO predicts cancer mortality will fall by about 5% by 2010 (from the 1995-7 baseline) but the gap in CHD mortality rates between Haringey and England will continue to increase.<sup>22</sup> There are significant inequalities across the borough with mortality rates from cancer in those under 75 in Northumberland Park in 2000-4 45% higher than the national average<sup>23</sup>.

### PRIMARY PREVENTION

See Sections on Smoking, Physical Activity, Food, Employment And Education

### SECONDARY PREVENTION

Action	Target group	Evidence of effectiveness	Delivery lead
Tackle low screening uptake rates for cervical and breast cancer including identification of communities that do not attend for screening, promotion of screening amongst low uptake groups, development of screening resources for non-English-speaking communities.	Women with low uptake of screening	Strong for certain interventions <sup>24</sup>	Screening co-ordinator

### TERTIARY PREVENTION (Treatment, Rehabilitation & Palliative Care)

Action	Target group	Evidence of effectiveness	Delivery lead
Implement and maintain cancer waiting times targets (time to see a specialist after GP referral, time to diagnosis, time to treatment)	Cancer patients	National Policy	HTPCT
Implementation of Integrated Cancer Care Programme	Cancer patients	Good practice	HTPCT Adult services
Extend the "Fit for Life" programme to cancer patients	Cancer patients	Good practice	HTPCT Public Health

## 6.ACCIDENTS

### Objective: (inc. PSA & local targets)

**PSA 5** Reduce the number of people killed or seriously injured in Great Britain in road accidents by 40% and the number of children killed or seriously injured by 50%, by 2010 compared with the average for 1994-98, tackling the significantly higher incidence in disadvantaged communities

### Current situation

Accidents are the leading cause of death in males under 20 in Haringey. As deaths from accidents occur at a relatively young age, they are the third most important cause of years of potential life lost (YPLL), after CVD and cancer. Land transport accidents account for nearly half of all deaths due to accidents. However, deaths and serious injuries caused by road traffic accidents have fallen from 131 in 2004 to 82 in 2005 and the gap between the borough and national average has been eliminated

Action	Target group	Evidence of effectiveness	Delivery lead
Maximise 20mph schemes and Safe Routes to School schemes	School children	Good practice	LBH Environmental Services
Ensure that accident prevention strategies are incorporate into home improvement schemes, particularly fire safety and prevention of trips and falls.	Households living in poor housing conditions.	Good practice	LBH Environmental Health
Development of local alcohol harm reduction strategy, inc. voluntary social responsibility scheme for alcohol retailers (code of practice and reporting of breaches), local authority enforcement, esp. sales to under 18s and alcohol screening and brief interventions in primary care and A&E	Will reflect strategy	Good practice, available, and evidence on a range of one-to-one interventions is expected.	DAAT
Maintain Children's Traffic Club for children aged 3+ to promote road safety.	Primary school children and parents	Good practice	Funded by Transport for London
Pilot alternative measures of traffic safety management- including Vehicle Activated Signs; priority give-ways; oversized mini-roundabouts; Homes Zones	To reflect intervention	Good practice	LBH Environmental Services



## 7.SUICIDE

### **Objective: (inc. PSA & local targets)**

Reduce mortality from suicide and undetermined injury by at least 20% by 2010.  
PSA05

### **Current situation**

The suicide mortality rate in Haringey has fallen from 10.7 per 100,000 population (10.7/100,000) in 1996/98 to 9.1/100,000 in 2002/04. If this trend continues, Haringey will meet the target 20% reduction by 2010. The gap between the Haringey and England average narrowed by 0.9/100,000 between 1996/98 and 2002/04 and is currently 0.4/100,000. Haringey had the third highest suicide mortality rate of its comparable boroughs in 2002/04, behind Lambeth (9.7/100,000) and Southwark (11.0/100,000). 75% of suicides in Haringey are amongst people who have not had contact with mental health services.

### **Action**

Continue to develop a Haringey suicide prevention strategy to include;

- Promotion of mental well-being amongst the wider population: building on findings from the Health in Mind project promoting access to support at early stages of mental distress through libraries and community settings.
- Reduction in the risk of suicide amongst key high-risk groups: including specific BME communities building on the 2006 report by Professor McKenzie.
- Reduction in the availability and lethality of suicide methods.

## 8.ACCESS TO HEALTH SERVICES

### Objective

Reduce number of Haringey residents not registered with a GP, and improve equity of access to health services.

### Current situation

There is little data on equity of access to services in Haringey. However, there is indirect evidence of inequity of access. In 2005, 955 Haringey residents had to be allocated a GP by the PCT, as they had approached 3 or more practices and been unable to register. The majority of these lived in the East of the borough. Despite CHD mortality being twice as high in some deprived wards in the east compared to more affluent boroughs in the west, standardised rates for CHD patients being treated in general practice and standardised hospital admission rates for CHD are not higher in the East of the borough, implying poor access to treatment.

Action	Target group	Evidence of effectiveness	Delivery lead
Work to develop one-stop-shops for health and social care services in accessible locations.	Service users	National policy	HTPCT, LBH, HSP
Equity audit of resource allocation to inform equitable commissioning of primary care services, and practice-based commissioning of services	Primary care population	Good practice	HTPCT-Commissioning Directorate
Improve funding and support for independent health advocates.	Vulnerable groups	Good practice	HTPCT teaching programme
Improve front-line health workers (e.g. receptionists) skills in communication and client care.	Service users	Good practice	HTPCT-Commissioning Directorate
Local enhanced service for the provision of services to patients who speak little or no English	Patients with little or no English	Good practice	HTPCT
Implement mental health enhanced service in primary care to improve/develop services that address the physical and mental health needs of people with mental health problems	Primary care service users with mental health problems	Good practice	HTPCT
Enhance involvement of voluntary sector and community groups in decision-making around service planning and development	Voluntary & community groups	Good practice	HTPCT, LBH, HAVCO & HARCEN
Improve transport services to hospitals/health services for disabled and older people	Disabled /older people	Good practice	HTPCT
Explore the role of libraries in providing information to inform health choices, and facilitating access to services.	Library service users	Good practice	LBH

## 9. INFANT MORTALITY

### Objective (inc. PSA and local targets)

Starting with children under one year, by 2010 reduce by at least 10% the gap in mortality between 'routine and manual' groups and the population as a whole.

PSA6a- Reducing the number of women who smoke during pregnancy

PSA6b- Increasing the number of women who initiate breastfeeding

### Current situation

The infant mortality rate in Haringey (7.4/1000 live births in 2002-2004) remains higher than London and England, and varies between Children's Network Area from 6.1/1000 in the West to 7.5 and 8.3 in the North and South patches respectively. Approximately 1 in 10 pregnant women in Haringey are current smokers at the time of delivery, twice the LDP target of 1 in 20. Approximately 84% of women in Haringey initiate breastfeeding, but data is not currently collected on breastfeeding maintenance. The Haringey Infant Mortality Action Plan 2004-5 is currently being reviewed, and this action plan will be updated in light of the outcomes.

Action	Target group	Evidence of effectiveness	Delivery lead
Develop a strategy to reduce the number of women booking late in their pregnancy for ante-natal care, in line with recent NICE guidance.	Pregnant women	Strong	Children's Service
Ensure new infant feeding coordinator role is able to promote breastfeeding and best practice in weaning, including implementation of infant feeding guidelines and development of programmes to promote breastfeeding that meet the Baby Friendly Initiative standards as a minimum.	Young children and parents/carers	Strong	Children's service
Systems to record and monitor the smoking status of, and interventions received by, families with children should be set up in line with NICE guidance. These systems should support service providers in providing smoking cessation support, for example at ante-natal appointments, delivery, during home visits, and other contacts.	Parents who smoke	Strong	Children's service
Smoking cessation services should be a core element of care pathways developed within children's centres.	Children's centre service users	Strong	Children's service
Develop a breastfeeding maintenance monitoring system using the child health surveillance system (6-8 week check), and use this to target interventions for women/families less likely to maintain breastfeeding.	Groups with low breastfeeding maintenance rates	Good practice	Children's Service

## 10.HOUSING

### Objective: (inc. PSA & local targets)

By 2010, bring all social housing into a decent condition with most of this improvement taking place in deprived areas, and for vulnerable households in the private sector, including families with children, increase the proportion who live in homes that are in decent condition (ODPM PSA7).

**LAA target:** tbc

### Current situation

Within the social housing sector, providers are on target to meet decent homes in 100% of stock by 2010.

The level of non-decent local authority owned housing stock has reduced from 58% in 2003/04 to 45% in March 2006. The majority of Registered Social Landlord (RSL) properties in Haringey meet the decent homes standard with approximately 80% of 10,500 properties meeting the standard as at April 2006

Action	Target group	Evidence of effectiveness	Delivery lead
Improve energy efficiency in private sector housing, especially homes which fail to meet standards due to a lack of thermal comfort.	Tenants in renewal areas	British Research Establishment modelling to identify key issues and areas for focus	LBH Environmental Health
Develop standard housing condition assessment criteria, guidance, and referral mechanisms to support services (eg private sector housing service) for a range of service providers visiting people in their own homes	Households living in poor accommodation that are vulnerable to poor health	Good practice	LBH Environmental Health
Implement system to ascertain and monitor levels of non-decency in the RSL sector.	Residents of non-decent housing	Good practice	LBH Housing Strategy
Implementation of Housing Association Forum joint service standards for all social landlords in Haringey.	Residents of social housing	Good practice	Housing Association Forum
Work with larger partner RSL associations and those which have more than 50% of properties failing to meet the Decent Homes standard, on their asset management plans to agree disposal programmes and with modified nominations agreements to enable decants for major works.	Tenants of larger RSLs failing to meet Decent Homes Standards	Good practice	LBH Housing Strategy

<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Delivery lead</b>
Implementation of Accredited Lettings Scheme to provide high quality private sector housing options	Tenants of private sector housing	Good practice	LBH Housing Strategy
Improve housing conditions in private rented sector accommodation above shops	Tenants of private sector housing above shops	Good practice	LBH Neighbourhood Management
Improve dilapidated private sector terrace properties in South Tottenham	Residents of private sector terrace properties in South Tottenham	Good practice	Bridge NDC
Develop initiatives to tackle fuel poverty	Residents living in fuel poverty	Strong evidence of links between fuel poverty and health outcomes	LBH Environmental Health
Continue to provide high quality floating support to those with housing support needs across all tenures through the supporting people programme	Residents with housing support needs	Good practice	LBH Supporting People Programme

## 11.EMPLOYMENT

### Objective: (inc. PSA & local targets)

**DWP PSA 4** In the 3 years to Spring 2008 demonstrate progress on increasing the employment rate; increase the employment rate of disadvantaged groups; significantly reduce the difference between the employment rate of disadvantaged groups and the overall rate.

**DWP PSA 8** In the three years to March 2008 increase the employment rate of disabled people, taking account of the economic cycle; and significantly reduce the difference between their employment rate and the overall rate, taking account of the economic cycle.

**DfES PSA 13** Increase the number of adults with the skills required for employability and progression to higher levels of training

**LAA target:** tbc

### Current situation

**Employment:** The employment rate amongst the total Haringey working age population was 60.3% in 2004/05. This was 14.5 percentage points below the England average of 74.8%. The gap between the Haringey and England average widened by 3.4 percentage points between 1997/98 and 2004/05, and is currently 14.5 percentage points.

**Education:** More than 85% of three-year-olds are accessing early years education. The attainment of 14 year-olds (Key Stage 3) has improved faster than the national trend since 2000, but the overall levels are still well below national figures. Although there is still a difference in attainment between schools in the East and West of Haringey, results in recent years suggest that this gap is also decreasing.

Action	Target group	Evidence of effectiveness	Delivery lead
Provide pre-employment training	Workless residents	Good practice	Urban futures
Contracts commissioned with Enfield College, Delta Club, John Grooms and Newton Housing delivering employment & skills support to Lone Parents, BME communities, Refugees, Disabled people	Lone Parents, BME communities, Refugees, Disabled people	Good practice	Enfield College Delta Club, John Grooms and Newton Housing
Provide pathways to work, flexible outreach services, generic and intensive support, job brokerage and work placements through the women stepping up programme	Women particularly from BME communities	Good practice	Haringey Women's Forum
Implement commissioned projects: <ul style="list-style-type: none"> <li>▪ Getting Haringey Working (At Work)</li> <li>▪ Employment Pathways to Health (Haringey Teaching PCT),</li> <li>▪ Learn for Work (I Can Do It Ltd.)</li> </ul>		Good practice	Haringey learning and skills partnership

<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Delivery lead</b>
KIS Business Challenge Assisting individuals in making the transition to self-employment by providing business start-up assistance to SMEs and young adults	SMEs and young adults	Good practice	
Continue to reduce the proportion of young people not in education, employment or training (NEET)	Young People	National policy	Connexions
Maximise Growth Area opportunities for new jobs and homes eg in Tottenham Hale, Hale Wharf and the London Stansted Cambridge Peterborough Corridor.	Residents in Tottenham Hale & Hale Wharf	Good practice	HSP
Continue to create new training opportunities to address the skills gap and get people into work.	Unemployed	Good practice	European Structural funds, LDA and Lottery funding
Health and Welfare to work Mental Health and employment project	People with mental health problems	Good practice	Richmond Fellowship
Workstep JCP contract to support disabled people gain and retain employment	Disabled people	Good practice	Haringey Council
Pilot effectiveness of offering employment advice in GP practices to target people on Incapacity Benefit who want to return to work.	People on incapacity benefit	Good practice	Haringey TPCT & Tomorrow's People (charity)

## 12. EDUCATION

### Objective: (inc. PSA & local targets)

**DfES PSA6** Raise standards in English and maths so that: y 2006, 85% of 11 year olds achieve level 4 or above, with this level of performance sustained to 2008; and by 2008, the proportion of schools in which fewer than 65% of pupils achieve level 4 or above is reduced by 40%.

**DfES PSA 7** Raise standards in English, maths, ICT and science in secondary education so that: by 2007, 85% of 14 year olds achieve level 5 or above in English, maths and ICT (80% in science) nationally, with this level of performance sustained to 2008; and by 2008, in all schools at least 50% of pupils achieve level 5 or above in each of English, maths and science.

**DfES PSA10** By 2008, 60% of those aged 16 to achieve the equivalent of 5 GCSEs at grades A\* to C; and in all schools at least 20% of pupils to achieve this standard by 2004, rising to 25% by 2006 and 30% by 2008.

**DfES PSA 13** Increase the number of adults with the skills required for employability and progression to higher levels of training

**LAA target:** tbc

### Current situation

Education: More than 85% of three-year-olds are accessing early years education. The attainment of 14 year-olds (Key Stage 3) has improved faster than the national trend since 2000, but the overall levels are still well below national figures. Although there is still a difference in attainment between schools in the East and West of Haringey, results in recent years suggest that this gap is also decreasing.

Action	Target group	Evidence of effectiveness	Delivery lead
Roll out of national EAL programme to improve English language competency for bilingual learners	Bilingual learners	Good practice	Children's Service
Support the introduction of Personal Advisors in 5 secondary schools to help pupils at risk of exclusion	Pupils at risk of exclusion	Good Practice	Children's Service
Development of programmes for secondary pupils from overseas who enter the education system at 14 plus. Programmes to ensure continuity into post 16 provision	Secondary pupils from overseas	Good practice	Children's service
Provide a wide range of Family Learning opportunities to parents and their children at pre-Foundation and Foundation Stage to boost early years attainment levels, particularly for those who are vulnerable.	Vulnerable pre-school children and parents	Good practice	CYPSP



<b>Action</b>	<b>Target group</b>	<b>Evidence of effectiveness</b>	<b>Delivery lead</b>
Support schools in developing provision that raises the achievement of Black and Minority Ethnic including promoting partnership between mainstream, supplementary and community language schools	BME children and young people	Good practice	CYPSP
Target schools where attendance is not improving consistently.	Children with poor school attendance	Good practice	CYPSP

## **Section 2: The case for action by the Haringey Strategic Partnership**

### **Introduction**

The purpose of the Haringey Life Expectancy Action Plan is to enable the Haringey Strategic Partnership to deliver priority actions to improve life expectancy and reduce health inequalities to meet the 2010 Public Service Agreement Targets.

### **National policy context**

Local authorities and primary care trusts have a responsibility for promoting the health and well being of their residents. Overall, people in Haringey are living longer healthier lives than they did 20 years ago. However, on average people in Haringey still die younger than in England as a whole, and there are substantial differences in health between neighbourhoods within the borough.

The causes of inequalities in health are multiple and complex. A small proportion of differences in health result from genetic and biological differences. The other influences on health are avoidable, and are the result of differences in:

- life circumstances (the opportunities we have in life, including our general socio-economic, cultural and environmental conditions);
- lifestyle (the choices we are able to make about how we live and their impact on health);
- access to services (our ability to have the same access to services whatever our background, age, or wherever we live).

Reducing disadvantage and health inequalities is a complex agenda that requires close partnership working across sectors and policy areas. This has been recognised by the Government in a number of policy initiatives over the past few years.

The 2003 report '*Tackling Health Inequalities: A Programme for Action*'<sup>25</sup> identified a key role for both national government and Local Strategic Partnerships in addressing the wider determinants of health inequalities.

The White Paper: '*Choosing Health; making healthier choices easier*'<sup>26</sup> emphasised the role of partnerships across communities, including local government, the NHS, business, the voluntary sector and faith communities in securing better access to healthier choices, especially for those in the most disadvantaged groups. '*Our health, our care, our say*' reiterated the importance of reducing health inequalities by improving access to health services, and through better prevention and earlier intervention.

### **What are the key targets that Haringey Strategic Partnership must meet?**

The Public Service Agreement targets of 2004 gave an increased profile to tackling inequalities in health. The targets aim to see faster improvements in health

outcomes amongst the 'fifth of areas with the worst health and deprivation indicators' in the country.

As Haringey falls in the bottom fifth of local authorities nationally for male and female life expectancy, heart and circulatory disease mortality and the Index of Multiple Deprivation (IMD) 2004 it has been designated one of the 88 'Spearhead LAs/PCTs'<sup>27</sup>.

As a member of the 'Spearhead' group, Haringey is aiming to meet the following Public Service Agreement Floor Targets by 2010:

- Reduce the gap in life expectancy by at least 10% between Haringey and the population as a whole
- Reduce mortality rates from heart disease and stroke and related diseases by at least 40% in people under 75, with at least a 40% reduction in the inequalities gap between Haringey and the population as a whole.
- Reduce mortality rates from cancer by at least 20% in people under 75, with a reduction in the inequalities gap between Haringey and the population as a whole of at least 6%
- Reduce mortality from suicide and undetermined injury by at least 20%
- Reduce the gap in infant mortality by at least 10% between "routine and manual groups" and the population as a whole
- Reduce adult smoking rates to 21% or less with a reduction in prevalence among routine and manual groups to 26% or less
- Halting the year-on-year rise in obesity among children under 11 by 2010 in the context of a broader strategy to tackle obesity in the population as a whole.
- Reduce the under -18 conception rate by 50% as part of a broader strategy to improve sexual health.

The full programme of Public Service Agreement Floor Targets includes a number of other targets which impact on health inequalities, including improvements in employment rates, housing, community safety, and education.

In addition, Haringey is negotiating local targets to address a number of local priorities through the Local Area Agreement (LAA) including;

- Narrowing the gap in premature mortality between Haringey and England, and between the most and least deprived wards in Haringey.
- Improving the uptake of smoking cessation services amongst people living in deprived areas
- Increasing physical activity amongst all ages, including older people

- Improving access to health services and homes for the most vulnerable
- Increasing the number of primary and secondary schools in the borough that meet the standards for Healthy School accreditation

### What is life expectancy?


Life expectancy is the number of years a baby born and living its whole life in an area would be expected to live if it were to experience the current (age-specific) death rates of that area. Life expectancy is best interpreted as a snapshot of the overall level of mortality in an area. It is not a forecast of how long babies will actually live, as current death rates are likely to change.<sup>28</sup> Nevertheless, it is a useful, easily understandable summary measure that can be used to compare death rates in different populations at different times. As deaths in earlier life contribute relatively more to lower life expectancy than deaths in older people, it also provides an indication of the number of premature deaths in an area.

Since age-specific death rates in men and women differ, life expectancy is usually calculated separately for each sex.

### What is the current life expectancy in Haringey?

The life expectancy for men and women in Haringey compared to London and England using mortality data from 1999-2003<sup>1</sup> is shown in figure 1. The lower life expectancy for men and women in Haringey compared to England and Wales is statistically significant<sup>2</sup>.

**Fig. 1 Life expectancy in Haringey compared to London and England, (pooled data from 1999-2003)**




<sup>1</sup> Combining data from several years helps to make the data more stable by reducing the influence of year-by-year variation in numbers of deaths.

<sup>2</sup> The error bars on the graph represent the 95% confidence intervals of the data. As the confidence intervals for the life expectancy in Haringey and London do not overlap, there is a 95% probability that the differences

## Is life expectancy in Haringey improving?

Along with national trends, life expectancy in Haringey for men and women has improved steadily over the past decade (see fig 2).

Fig 2. Trends in Life Expectancy for Haringey and England and Wales (E&W) 1991-2003



Due to year on year fluctuations in mortality rates at the small area level, it is not possible to use current trends to predict whether the life expectancy gap between Haringey and England as a whole is likely to widen or narrow by 2010. However, at both the London level<sup>29</sup> and the national level<sup>30</sup> the gap in life expectancy at birth between England and the Spearhead Group continues to widen. Therefore it is likely that the gap between Haringey and England will widen unless specific action is taken to improve the health of the most disadvantaged groups.

## Does life expectancy vary within Haringey?

Within Haringey, life expectancy varies significantly between different wards. The variation in life expectancy between wards in Haringey is even greater than the variation in life expectancy between local authorities in London<sup>31</sup>.

---

between the figures for Haringey and London are real and not due to chance year-by-year variations in death rates.

Figure 3 shows the variation in male life expectancy between wards in Haringey. Generally, the more deprived wards (as measured by the Index of Multiple Deprivation 2004) have a lower male life expectancy than the more affluent wards. At the two extremes, male life expectancy in Bruce Grove (70.5 years) is nearly 8 years lower than male life expectancy in Muswell Hill (78.2 years). The relationship between male life expectancy and ward-level deprivation is strong and statistically significant.

**Figure 3. Male life expectancy 1999-2003 by ward in Haringey**





Figure 4 shows the variation in female life expectancy between wards in Haringey. There is only a weak relationship between female life expectancy and deprivation, and this is not statistically significant.

**Fig 4. Female life expectancy 1999-2003 by ward in Haringey**




A stronger relationship between life expectancy and deprivation for men than for women is also found across London<sup>32</sup> and at the national level<sup>33</sup>. The reasons for this are not fully understood. Previous studies have speculated that this might be due to a stronger association between deprivation and health risk behaviours in men than women, or because men with poor health may be more likely to migrate to more deprived areas.

## What causes of early death impact most on life expectancy in Haringey?

Figure 5 shows the main causes of premature death (deaths under the age of 75 years) in Haringey over the 3-year period from 2001-2003.

Fig 5. Main causes of death for persons <75 years in Haringey 2001-2003 (numbers and percent)



As shown, heart and circulatory diseases and cancer together account for 67% of all premature deaths in Haringey.

Deaths occurring earlier in life contribute relatively more to lower life expectancy than deaths in later life. One way of looking at the causes of death that contribute most to life expectancy is by calculating, for each cause of death, the number of years that people would have lived had they lived until they were 75. This is known as the Years of Potential Life Lost (YPLL).



Table 1 shows that heart and circulatory diseases and cancer account for around half of all the years of potential life lost. However, accidents and suicide, and injuries of undetermined intent also account for a significant proportion of YPLL (20% in males and 9% in females). This is because these causes of death disproportionately affect younger people, and so contribute more to years of potential life lost and life expectancy than to overall mortality rates.

**Table 1. Main causes of Years of Potential Life Lost (YPLL) in Haringey 2001-3**

Cause	Males – number of YPLL (%)	Females - number of YPLL (%)
<b>All heart and circulatory diseases</b>	4,853 (25)	2,579 (22)
<b>All cancers</b>	4,279 (22)	3,911 (33)
<b>Accidents</b>	2,317 (12)	390 (3)
<b>Suicide and injuries of undetermined intent</b>	1,617 (8)	692 (6)
<b>Infectious and parasitic disease</b>	805 (4)	433 (4)
<b>Respiratory disease</b>	596 (3)	635 (6)

### **How are the main causes of premature death distributed in Haringey?**

To compare the distribution of deaths between different populations it is important to take into account not just the number of deaths, but also the size of the populations and their age profiles. The commonest way to do this is by calculating the Standardised Mortality Ratio (SMR)<sup>3</sup>.

---

<sup>3</sup> The SMR is the ratio of the number of deaths occurring in a population to the number that would have occurred if that population had the same age-specific death rates as the population of England and Wales. The ratio is multiplied by 100. An SMR of 100 means that a population has the same age-specific death rates as the England and Wales population. An SMR of 120 means that a population has 20% more age-specific deaths than the E&W population. An SMR of 80 means that a population has a 20% lower age-specific death rate than the E&W population.

Figure 6 shows the Standardised Mortality Ratio for Coronary Heart Disease (the most common cause of death due to heart and circulatory disease) for persons under the age of 75 by ward. Northumberland Park and Bruce Grove (the most deprived wards in Haringey as measured by IMD 2004) have mortality rates due to Coronary Heart Disease (CHD) more than 70% higher than the average CHD mortality rates in England and Wales. There is a statistically significant relationship between SMR for coronary heart disease and ward-level deprivation in Haringey.

Figure 6. Standardised Mortality Ratio for Coronary Heart Disease by ward in Haringey, 2000-2004





Figure 7 shows the Standardised Mortality Ratio for cancer for persons aged under 75 years by ward. Again, there is a statistically significant relationship between SMR for cancer and ward-level deprivation in Haringey.

Figure 7 Standardised Mortality Ratio for Cancer by ward in Haringey, 2000-2004



## **What are the main determinants of inequalities in life expectancy in Haringey?**

As mentioned earlier, the causes of inequalities in health are complex and relate to a combination of people's social and economic circumstances, their access to services and their personal behaviour, which is itself influenced by the social and cultural environment. However, there are a number of clear risk factors for the main causes of premature death and inequalities in health in Haringey that are amenable to change:

- **Smoking**
  - Smoking is the individual health behaviour with the single largest impact on health inequalities.
  - Smoking is a major risk factor for heart and circulatory diseases, lung cancer, chronic lung disease and many other conditions.
  - The prevalence of smoking is considerably higher amongst people of lower socio-economic class, lone parents, the unemployed and people with mental illness than amongst the rest of the population<sup>34</sup>.
  - It has been estimated that around two thirds of the observed difference in risk of death across social groups in middle age is caused by smoking tobacco<sup>35</sup>.
  - Reducing smoking will result in substantial reductions in mortality from coronary heart disease within 12-24 months<sup>36</sup>
  
- **Food and nutrition**
  - High blood pressure (which is directly related to obesity and high salt intake) and high serum cholesterol (which is directly linked to high intakes of saturated fat) are the two main risk factors for diseases of the heart and circulatory system<sup>37</sup>.
  - Low fruit and vegetable intake is closely linked with a high prevalence of some cancers and heart and circulatory disease.
  - Poorer households in poorer communities are less likely to have access to healthy, affordable food.
  - Poorer households eat less fruit and vegetables, salad, wholemeal bread, wholegrain and high-fibre cereals and oily fish, and more white bread, full-fat milk, table sugar and processed meat products.
  
- **Physical activity**
  - People who have a physically active lifestyle are at approximately half the risk of developing heart disease compared to those who have a sedentary lifestyle<sup>38</sup>.
  - Regular physical activity is also associated with a reduced risk of diabetes, obesity, osteoporosis and colon cancer, and with improved mental health.
  - In older adults physical activity is associated with increased functional capacities.
  - Physical inactivity is associated with low social class, income and educational attainment, indicating that developing opportunities for physical activity is particularly important in these groups
  
- **Housing**

- Housing affects people's physical and mental health in a range of ways, from the quality of the indoor environment to neighbourhood quality and safety and housing allocation and homelessness<sup>39</sup>.
  - In Haringey a significant proportion of local authority homes and private rented homes are considered to be non-decent.
  - The most vulnerable people live in non-decent homes: people who live alone, ethnic minorities and households with no one in full-time employment are most likely to live in such accommodation.
- **Employment**
    - Employment status is a key determinant of income and social status, and thus closely linked with health and health inequalities.
    - A middle-aged man who loses his job is twice as likely to die in the next 5 years as a man who remains in employment.
    - Worklessness and workless households are highly concentrated in particular neighbourhoods. This has important implications for community regeneration and the economic vitality of neighbourhoods.
- **Education**
    - Education influences health in a variety of ways.
    - Educational qualifications are an important determinant of employment prospects, which in turn influence access to income and material resources.
    - Education also provides children and young people with the knowledge and skills to lead a healthier life
    - The educational attainment of 14-year olds and 16-year olds in Haringey schools are well below the national average. However, attainment in Haringey schools is improving faster than the national average, and the gap between schools in the east and the west of the borough is closing
- **Accidents**
    - Accidents were the leading cause of death in under 20 year olds in Haringey in 2001-2
    - Accidental death is much more common amongst males than females.
    - Road traffic accidents account for more than half of accidental deaths in Haringey.
    - Local data show that more than a quarter of child pedestrian casualties happen in the 10% most deprived wards.
- **Suicide**
    - Suicide is a significant contributor to early death in Haringey.
    - In Haringey, approximately 35 people commit suicide in 2001, which is more than 50% higher than the national average. This is in part due to the high levels of factors increasing the risk of suicide, such as mental illness, unemployment, substance misuse and social exclusion.
    - Three quarters of suicides in Haringey are amongst people who have not had contact with mental health services

- **Health services**

- There are a number of health service interventions that can significantly reduce mortality amongst patients with heart disease and cancer and those at high risk for these diseases. Most important are those that reduce risk factors for the development of heart disease (smoking cessation services, treatment of hypertension and the use of statins to reduce the risk of cardio-vascular events in those at risk of heart disease or with established heart disease) and the early detection and treatment of cancers.
- The 2010 time-scale for the life expectancy, cancer and heart disease targets means that we need to focus attention on reducing premature death amongst those that already have, or are at high risk of developing these diseases<sup>40</sup>.
- There are a number of barriers to accessing good quality health services, and there is evidence that those who are most vulnerable often have poorest access to services.

## References

<sup>1</sup> A Healthier Haringey: Improving wellbeing and tackling inequalities, report of an event on 8th February 2006'

<sup>3</sup> HDA. Smoking and health inequalities.2002

<sup>4</sup> Scollo M et al, Review of the Quality of Studies on the Economic Effects of Smoke-free Policies on the Hospitality Industry. Tobacco Control 2003;12:13-20.

<sup>5</sup> NICE Smoking cessation guidance. March 2006. <http://www.nice.org.uk/page.aspx?o=299611>

<sup>6</sup> London Health Observatory. Stop before the Op. May 2006.

<http://www.lho.org.uk/Download/fqb2zui11ml0hb2u2hg5z42h/live/10495/Stop%20before%20the%20Op%20Final.pdf>

<sup>7</sup> Department of Health (2004) Choosing Health. Making healthier choices easier

<sup>8</sup> Department of Health (1999) Health Survey for England 2003

<sup>9</sup> Britton, A, McPherson, K. (2000) Monitoring the progress of the 2010 target for coronary heart disease mortality: Estimated consequences on CHD incidence and mortality from changing prevalence of risk factors. National Heart Forum: London

<sup>10</sup> NICE Physical activity guidance. March 2006. <http://www.nice.org.uk/page.aspx?o=299531>

<sup>11</sup> Breastfeeding reference

<sup>12</sup> LHO Mortality from all circulatory diseases 2002-4

<sup>13</sup> LHO Standardised mortality ratio CHD 2000-4

<sup>14</sup> LHO The London Forecast. Can London's health divide be reduced? 2004

<sup>15</sup> DH Quality and Outcomes Framework (QOF) guidance and evidence base 2004.

<http://www.dh.gov.uk/assetRoot/04/08/86/93/04088693.pdf>

<sup>16</sup> NICE. Statins for the prevention of cardiovascular events. Jan 2006.

<http://www.nice.org.uk/page.aspx?o=TA094guidance>

<sup>17</sup> DH Quality and Outcomes Framework (QOF) guidance and evidence base 2004.

<http://www.dh.gov.uk/assetRoot/04/08/86/93/04088693.pdf>

<sup>18</sup> NICE Management of chronic heart failure in adults in primary and secondary care 2003

<http://www.nice.org.uk/page.aspx?o=CG005>

<sup>19</sup> NICE guideline on prophylaxis for patients who have experienced a myocardial infarction 2001

<http://www.nice.org.uk/page.aspx?o=16529>

<sup>20</sup> NICE The clinical effectiveness and cost effectiveness of early thrombolysis for treatment of myocardial infarction 2002 <http://www.nice.org.uk/page.aspx?o=ta052&c=cardiovascular>

<sup>21</sup> LHO Standardised mortality ratio cancer 2000-2004

<sup>22</sup> LHO The London Forecast. Can London's health divide be reduced? 2004

<sup>23</sup> LHO Standardised mortality ratio cancer 2000-2004

<sup>24</sup> HTA The determinants of screening uptake and interventions for increasing uptake: a systematic review. 2000.

<http://www.hta.nhsweb.nhs.uk/fullmono/mon414.pdf>

<sup>25</sup> Department of Health. Tackling Health Inequalities: a programme for action. 2003.

<http://www.dh.gov.uk/assetRoot/04/01/93/62/04019362.pdf>

- 
- <sup>26</sup> Department of Health. Choosing Health: making healthier choices easier. 2004.  
[http://www.dh.gov.uk/PublicationsAndStatistics/Publications/PublicationsPolicyAndGuidance/PublicationsPolicyAndGuidanceArticle/fs/en?CONTENT\\_ID=4094559&chk=H29Li6](http://www.dh.gov.uk/PublicationsAndStatistics/Publications/PublicationsPolicyAndGuidance/PublicationsPolicyAndGuidanceArticle/fs/en?CONTENT_ID=4094559&chk=H29Li6)
- <sup>27</sup> The Healthcare Standards Unit. Tackling health inequalities. The Spearhead group of local authorities. 2005.  
[www.hcsu.org.uk/index.php?option=com\\_docman&task=doc\\_download&gid=271](http://www.hcsu.org.uk/index.php?option=com_docman&task=doc_download&gid=271)
- <sup>28</sup> Fitzpatrick J, Jacobson B. Mapping health inequalities across London. LHO, September 2001. Calculating life expectancy and infant mortality rates. Mapping health inequalities across London: technical supplement.
- <sup>29</sup> London Health Observatory. The London Health Forecast: Can London's health divide be reduced? 2004.  
<http://www.lho.org.uk/viewResource.aspx?id=8990>
- <sup>30</sup> Department of Health. Tackling health inequalities: status report on the programme for action. August 2005  
<http://www.dh.gov.uk/assetRoot/04/11/76/98/04117698.pdf>
- <sup>31</sup> Health inequalities in London. Life expectancy and infant mortality 1998-2002. London Health Observatory, April 2004
- <sup>32</sup> London Health Observatory. Health inequalities in London: Life expectancy and mortality. 2004
- <sup>33</sup> London Health Observatory. Mapping health inequalities across London. 2001.  
[http://www.lho.org.uk/Download/n3eq2c55x2pvaynmzdhwr0j5/live/7652/map\\_hilond\\_3.pdf](http://www.lho.org.uk/Download/n3eq2c55x2pvaynmzdhwr0j5/live/7652/map_hilond_3.pdf)
- <sup>34</sup> Social patterning of individual health behaviours: the case of cigarette smoking. Jarvis MJ and Wardle J. In Marmot M and Wilkinson RG. Social Determinants of Health. Oxford. Oxford University Press. 2006
- <sup>35</sup> Social patterning of individual health behaviours: the case of cigarette smoking. Jarvis MJ and Wardle J. In Marmot M and Wilkinson RG. Social Determinants of Health. Oxford. Oxford University Press. 2006.
- <sup>36</sup> Kelly M and Capewell S 2004. Relative contributions of changes in risk factors and treatment to the reduction in coronary heart disease mortality. Health Development Agency Briefing Paper<sup>36</sup>
- <sup>37</sup> World Health Organisation 2004. Food and health in Europe: a new basis for action.  
<http://www.euro.who.int/document/e78578.pdf>
- <sup>38</sup> Health development Agency 2004. The effectiveness of public health interventions for increasing physical activity among adults evidence briefing <http://www.publichealth.nice.org.uk/download.aspx?o=502697>
- <sup>39</sup> National Institute for Health and Clinical Excellence. 2005. Housing and public health: a review of reviews of interventions for improving health - Evidence briefing  
<http://www.publichealth.nice.org.uk/page.aspx?o=526671>
- <sup>40</sup> Department of Health 2005. Tackling health inequalities: what works?.  
<http://www.dh.gov.uk/assetRoot/04/10/34/06/04103406.pdf>