

PRIMARY CARE**Quality and Outcomes framework 2007/08, produced per kind favour by Mahnaz Shaukat, Public Health Strategist, Islington PCT**

The following data has been shared by the public health intelligence leads in each PCT, and put together by Islington PCT. Please note that the data has been extracted from the QMAS system as at the end of March 2008 and may be subject to change when published nationally in June.

NB: 4 Camden practices were excluded from the minimum, maximum score as these practices have very specific populations (homeless, students etc) and hence skew the range.

Prevalence

Table 1: Number of patients on the CHD, Stroke and Heart Failure register and crude prevalence using GP list size as of February 2008.

Register	Islington Crude prevalence % (number on register)	Camden Crude prevalence % (number on register)	Enfield Crude prevalence % (number on register)	Haringey Crude prevalence % (number on register)	Barnet Crude prevalence % (number on register)
CHD	2.0% (3,913)	1.97% (4,519)	2.58% (7,555)	2.09% (5,731)	2.8% (10,338)
Stroke	1.1% (2,169)	0.98% (2,252)	1.09% (3,197)	0.84% (2,317)	0.94% (3,428)
Heart Failure	0.60% (1,202)	0.50% (1,152)	0.54% (1,589)	0.45% (1,245)	0.59% (2,167)

Table 2: Average percentage achievement for CHD indicators and range of achievement by practice.

QOF 2007/08		Islington	Camden	Enfield	Haringey	Barnet
CHD 2	Patients with newly diagnosed angina	95% (40%-100%)	93% (60%-100%)	90% (0%-100%)	80% (6%-100%)	94% 95%-100%)
CHD 5	Record of blood pressure in the last 15 months	98% (94%-100%)	97% (88%-100%)	98% (68%-100%)	97% (82%-100%)	97% (80%-100%)
CHD 6	Last blood pressure record is <150/90	91% (78%-100%)	86% (73%-100%)	90% (63%-100%)	88% (55%-100%)	89% (71%-100%)
CHD 7	Record of cholesterol in the last 15 months	94% (86%-98%)	91% (57%-100%)	92% (33%-100%)	91% (48%-100%)	93% (51%-100%)
CHD 8	Last cholesterol is <5.0ml	83% (69%-97%)	79% (42%-96%)	78% (26%-96%)	78% (39%-100%)	80% (48%-98%)
CHD 9	CHD patients on antiplatelet therapy/anticoagulation	95% (88%-100%)	93% (68%-100%)	95% (74%-100%)	93% (71%-100%)	94% (71%-100%)
CHD 10	CHD patients treated with a beta blocker	75% (37%-100%)	70% (30%-90%)	73% (30%-100%)	70% (36%-100%)	68% (41%-100%)
CHD 11	Patients with a history of MI who are currently treated with ACE inhibitor	94% (80%-100%)	88% (50%-100%)	91% (57%-100%)	90% (33%-100%)	88% (71%-100%)

Table 3: Average percentage achievement for Heart Failure indicators and range of achievement by practice.

QOF 2007/08		Islington	Camden	Enfield	Haringey	Barnet
HF 2	Patients with a diagnosis of heart failure confirmed with ECHO	98% (75%-100%)	96% (75%-100%)	89% (0%-100%)	94% (0%-100%)	96% (0%-100%)
HF3	Patients with a current diagnosis of heart failure due to LVD who are currently treated with an ACE inhibitor or Angiotensin Receptor Blocker, who can tolerate therapy and for whom there is no contra-indication.	92% (80%-100%)	89% (67%-100%)	90% (0%-100%)	90% (0%-100%)	90% (0%-100%)

Table 4: Average percentage achievement for Stroke indicators and range of achievement by practice.

QOF 2007/08		Islington	Camden	Enfield	Haringey	Barnet
Stroke 5	Record of Blood Pressure in last 15 months	98% (91%-100%)	96% (75%-100%)	98% (72%-100%)	96% (67%-100%)	96% (80%-100%)
Stroke 6	Last BP is < 150/90	90% (74%-100%)	85% (72%-100%)	89% (67%-100%)	86% (57% - 100%)	88% (70%-100%)
Stroke 7	Record of cholesterol check in last 15 months	92% (76%-100%)	89% (65%-100%)	89% (50%-100%)	87% (33% -100%)	90% (33%-100%)
Stroke 8	Last record of cholesterol is <5.0m/ml	77% (65%-100%)	75% (54%-94%)	72% (25%-92%)	71% (25% -100%)	74% (33%-100%)
Stroke 11	New stroke patients referred for further investigation	92% (0%-100%)	91% (67%-100%)	84% (0%-100%)	88% (0% -100%)	93% (50%-100%)
Stroke 12	TIA/Ischaemic Stroke on antiplatelet/anticoagulation	95% (80%-100%)	93% (75%-100%)	95% (75%-100%)	94% (57% -100%)	94% (75%-100%)